

Clinical Policy: Lanadelumab-fylo (Takhzyro)

Reference Number: CP.PHAR.396

Effective Date: 12.01.18

Last Review Date: 02.22

Line of Business: Commercial, HIM, Medicaid

[Coding Implications](#)

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

Description

Lanadelumab-fylo (Takhzyro[™]) is a human monoclonal antibody that inhibits the proteolytic activity of kallikrein to reduce the generation of bradykinin.

FDA Approved Indication(s)

Takhzyro is indicated for prophylaxis to prevent attacks of hereditary angioedema (HAE) in patients 12 years and older.

Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of health plans affiliated with Centene Corporation[®] that Takhzyro is **medically necessary** when the following criteria are met:

I. Initial Approval Criteria

A. Hereditary Angioedema (must meet all):

1. Diagnosis of HAE confirmed by a history of recurrent angioedema and one of the following (a or b):
 - a. Low C4 level and low C1-INH antigenic or functional level (*see Appendix D*);
 - b. Normal C4 level and normal C1-INH level, and at least one of the following (i or ii):
 - i. Presence of a mutation associated with the disease (*see Appendix D*);
 - ii. Family history of angioedema and documented failure of high-dose antihistamine therapy (i.e., cetirizine 40 mg/day or equivalent) for at least 1 month or an interval expected to be associated with 3 or more attacks of angioedema, whichever is longer;
2. Prescribed by or in consultation with an allergist, hematologist, or immunologist;
3. Age \geq 12 years;
4. Prescribed for long-term prophylaxis of HAE attacks and request meets one of the following (a, b, or c):
 - a. Member experiences more than one severe event per month;
 - b. Member is disabled more than five days per month;
 - c. Member has history of previous airway compromise;
5. Failure of Haegarda[®], unless contraindicated or clinically significant adverse effects are experienced;

6. Member is not using Takhzyro in combination with another FDA-approved product for long-term prophylaxis of HAE attacks (e.g., Cinryze[®], Haegarda[®], Orladeyo[™]);
7. Dose does not exceed 300 mg every 2 weeks.

Approval duration:

Medicaid/HIM – 6 months

Commercial – 6 months or to the member’s renewal date, whichever is longer

B. Other diagnoses/indications

1. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid.

II. Continued Therapy

A. Hereditary Angioedema (must meet all):

1. Currently receiving medication via Centene benefit or member has previously met initial approval criteria;
2. Member is responding positively to therapy as evidenced by reduction in attacks from baseline;
3. Member is not using Takhzyro in combination with another FDA-approved product for long-term prophylaxis of HAE attacks (e.g., Cinryze, Haegarda, Orladeyo);
4. Request is for 300 mg every 4 weeks, unless documentation supports member is not well-controlled (e.g., attack(s) within the last 6 months);
5. If request is for a dose increase, new dose does not exceed 300 mg every 2 weeks.

Approval duration:

Medicaid/HIM – 12 months

Commercial – 6 months or to the member’s renewal date, whichever is longer

B. Other diagnoses/indications (must meet 1 or 2):

1. Currently receiving medication via Centene benefit and documentation supports positive response to therapy.

Approval duration: Duration of request or 6 months (whichever is less); or

2. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid.

III. Diagnoses/Indications for which coverage is NOT authorized:

- A.** Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid or evidence of coverage documents.

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

CI-INH: C1 esterase inhibitor

C4: complement component 4

FDA: Food and Drug Administration
HAE: Hereditary Angioedema

HAE-nl-C1INH: hereditary angioedema
with normal C1 inhibitor

Appendix B: Therapeutic Alternatives

This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent for all relevant lines of business and may require prior authorization.

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
cetirizine	40 mg/day (<i>off-label</i>) Typical dosing range (mg/day): 10 mg/day <i>US HAEA Medical Advisory Board 2020 Guidelines for the Management of Hereditary Angioedema</i>	40 mg/day (<i>off-label</i>)
C1 esterase inhibitor (Haegarda [®])	60 IU/kg body weight SC twice weekly (every 3 or 4 days)	Based on weight, 60 IU/kg/dose

Therapeutic alternatives are listed as Brand name[®] (generic) when the drug is available by brand name only and generic (Brand name[®]) when the drug is available by both brand and generic.

Appendix C: Contraindications/Boxed Warnings

None reported

Appendix D: General Information

- Diagnosis of HAE:
 - There are two classifications of HAE: HAE with C1-INH deficiency (HAE-C1INH, further broken down into Type I and Type II) and HAE with normal C1-INH (also known as HAE-nl-C1INH). HAE-nl-C1INH was previously referred to as type III HAE, but this term is obsolete and should not be used.
 - In both Type I (~85% of cases) and Type II (~15% of cases), C4 levels are low. C1-INH antigenic levels are low in Type I while C1-INH functional levels are low in Type II. Diagnosis of Type I and II can be confirmed with laboratory tests. Reference ranges for C4 and C1-INH levels can vary across laboratories (see below for examples); low values confirming diagnosis are those which are below the lower end of normal.

Laboratory Test & Reference Range	Mayo Clinic	Quest Diagnostics	Lab Corp
C4	14 – 40 mg/dL	13-57 mg/dL (age- and gender-specific ranges)	14 – 44 mg/dL
C1-INH, antigenic	19 – 37 mg/dL	21 – 39 mg/dL	21 – 39 mg/dL
C1-INH, functional	Normal: > 67% Equivocal: 41 – 67% Abnormal: < 41%	Normal: ≥ 68% Equivocal: 41 – 67% Abnormal: ≤ 40%	Normal: > 67% Equivocal: 41 – 67% Abnormal: < 41%

- HAE-nl-C1INH, on the other hand, presents with normal C4 and C1-INH levels. Some patients have a known associated mutation, while others have no identified genetic indicators. HAE-nl-C1INH is very rare, and there are no laboratory tests to confirm the diagnosis; mutations in 4 genes causing HAE-nl-C1INH have been identified:

Identified Genes Associated with Mutations in HAE-nl-C1INH
<i>F12</i>
<i>ANGPT1</i>
<i>PLG</i>
<i>KNG1</i>

V. Dosage and Administration

Indication	Dosing Regimen	Maximum Dose
HAE attack prophylaxis	300 mg SC every 2 weeks A dosing interval of 300 mg every 4 weeks may be considered if the patient is well-controlled (e.g., attack free) for more than 6 months	300 mg SC every 2 weeks

VI. Product Availability

Injections: 300 mg/2 mL (150 mg/mL) solution in single-dose vial and single-dose prefilled syringe

VII. References

1. Takhzyro Prescribing Information. Lexington, MA: Shire ViroPharma Incorporated; February 2022. Available at: <https://www.Takhzyro.com/>. Accessed March 1, 2022.
2. Maurer M, Mager M, Ansotegui I, et al. The International WAO/ESSCI guideline for the management of hereditary angioedema – the 2017 revision and update. *World Allergy Organ J*. 2018; 11:5
3. Cicardi M, Aberer W, Banerji A, et al. Classification, diagnosis, and approach to treatment for angioedema: consensus report from the Hereditary Angioedema International Working Group. *Allergy*. 2014; 69(5): 602-616.
4. Zuraw B, Bernstein J, Lang D. A focused parameter update: Hereditary angioedema, acquired C1 inhibitor deficiency, and angiotensin-converting enzyme inhibitor-associated angioedema. *J Allergy Clin Immunol*. 2013; 131(6): 1491-3.
5. Busse PJ, Christiansen SC, Reidl MA, et al. US HAEA Medical Advisory Board 2020 Guidelines for the Management of Hereditary Angioedema. *J Allergy Clin Immunol*. 2021; 9(1): 132-150.e3.
6. Mayo Clinic Laboratories [internet database]. Rochester, Minnesota: Mayo Foundation for Medical Education and Research. Updated periodically. Accessed November 8, 2021.
7. Quest Diagnostics ® [internet database]. Updated periodically. Accessed November 8, 2021.
8. LabCorp [internet database]. Burlington, North Carolina: Laboratory Corporation of America. Updated periodically. Accessed November 8, 2021.

Coding Implications

Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

HCPCS Codes	Description
J0593	Injection, lanadelumab-flyo, 1 mg

Reviews, Revisions, and Approvals	Date	P&T Approval Date
Policy created	09.25.18	11.18
1Q 2019 annual review: added requirement that member is not using requested product in combination with other approved products for the long-term prophylaxis of HAE attacks; references reviewed and updated.	11.19.18	02.19
Added HIM line of business per SDC and prior approved clinical guidance.	04.01.19	
1Q 2020 annual review: HAE lab reference range updated; removed rheumatologist specialty for alignment; revised dosing criteria for dose reduction if member is well-controlled per PI; added coding implications; references reviewed and updated.	11.04.19	02.20
1Q 2021 annual review: no significant changes; references reviewed and updated.	10.08.20	02.21
Per June SDC and prior clinical guidance, added redirection to Haegarda.	06.02.21	08.21
1Q 2022 annual review: updated diagnosis criteria to include a recurrent history of angioedema and either an associated mutation or family history of angioedema with failure of high-dose antihistamines for HAE-nl-C1INH; references reviewed and updated.	11.08.21	02.22
RT4: New dose formulation of prefilled syringe added to policy.	03.01.22	

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. “Health Plan” means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan’s affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

Note:

For Medicaid members, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

©2018 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or

remove any trademark, copyright or other notice contained herein. Centene[®] and Centene Corporation[®] are registered trademarks exclusively owned by Centene Corporation.