

## **Clinical Policy: Bempedoic Acid (Nexletol), Bempedoic Acid/Ezetimibe (Nexlizet)**

Reference Number: CP.PMN.237

Effective Date: 09.01.20

Last Review Date: 02.22

Line of Business: Commercial, HIM, Medicaid

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

### **Description**

The following are adenosine triphosphate-citrate lyase (ACL) inhibitors requiring prior authorization: bempedoic acid (Nexletol<sup>™</sup>) and bempedoic acid/ezetimibe (Nexlizet<sup>™</sup>). Nexlizet contains ezetimibe, which is a cholesterol absorption inhibitor.

### **FDA Approved Indication(s)**

Nexletol and Nexlizet are indicated for use as adjuncts to diet and maximally tolerated statin therapy for the treatment of adults with heterozygous familial hypercholesterolemia (HeFH) or established atherosclerotic cardiovascular disease (ASCVD) who require additional lowering of low-density lipoprotein cholesterol (LDL-C).

Limitation(s) of use: The effect of Nexletol and Nexlizet on cardiovascular morbidity and mortality has not been determined.

### **Policy/Criteria**

*Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.*

It is the policy of health plans affiliated with Centene Corporation<sup>®</sup> that Nexletol and Nexlizet are **medically necessary** when the following criteria are met:

#### **I. Initial Approval Criteria**

##### **A. Heterozygous Familial Hypercholesterolemia and Atherosclerotic Cardiovascular Disease (must meet all):**

1. Diagnosis of one of the following (a or b):
  - a. ASCVD as evidenced by a history of any one of the following conditions (i-vii):
    - i. Acute coronary syndromes;
    - ii. Clinically significant coronary heart disease (CHD) diagnosed by invasive or noninvasive testing (such as coronary angiography, stress test using treadmill, stress echocardiography, or nuclear imaging);
    - iii. Coronary or other arterial revascularization;
    - iv. Myocardial infarction;
    - v. Peripheral arterial disease presumed to be of atherosclerotic origin;
    - vi. Stable or unstable angina;
    - vii. Stroke or transient ischemic attack (TIA);
  - b. HeFH, and member meets both of the following (i and ii):

- i. Baseline LDL-C (prior to any lipid-lowering pharmacologic therapy) was  $\geq$  190 mg/dL;
  - ii. HeFH diagnosis is confirmed by one of the following (a or b):
    - a) World Health Organization (WHO)/Dutch Lipid Network familial hypercholesterolemia diagnostic criteria score of  $> 8$  as determined by requesting provider (*see Appendix D*);
    - b) Definite diagnosis per Simon Broome criteria (*see Appendix D*);
2. Prescribed by or in consultation with a cardiologist, endocrinologist, or lipid specialist;
3. Age  $\geq$  18 years;
4. For members on statin therapy, both of the following (a and b):
  - a. Nexletol or Nexlizet is prescribed in conjunction with a statin at the maximally tolerated dose;
  - b. Member has been adherent for at least the last 4 months to maximally tolerated doses of one of the following statin regimens (i, ii, or iii):
    - i. A high intensity statin (*see Appendix E*);
    - ii. A moderate intensity statin (*see Appendix E*), and member has one of the following (a or b):
      - a) Intolerance to two high intensity statins;
      - b) A statin risk factor (*see Appendix G*);
    - iii. A low intensity statin and member has one of the following (a or b):
      - a) Intolerance to one high and one moderate intensity statins;
      - b) A statin risk factor (*see Appendix G*) and history of intolerance to two moderate intensity statins;
5. For members not on statin therapy, member meets one of the following (a or b):
  - a. Statin therapy is contraindicated per Appendix F;
  - b. For members who are statin intolerant, member has tried at least two statins, one of which must be hydrophilic (pravastatin, fluvastatin, or rosuvastatin), and member meets one of the following (i or ii):
    - i. Member has documented statin risk factors (*see Appendix G*);
    - ii. Member is statin intolerant due to statin-associated muscle symptoms (SAMS) and meets both of the following (a and b):
      - a) Documentation of intolerable SAMS persisting at least two weeks, which disappeared with discontinuing the statin therapy and recurred with a statin re-challenge;
      - b) Documentation of re-challenge with titration from lowest possible dose and/or intermittent dosing frequency (e.g., 1 to 3 times weekly);
6. Member has been adherent to ezetimibe therapy used concomitantly with a statin at the maximally tolerated dose for at least the last 4 months, unless contraindicated per Appendix F or member has a history of ezetimibe intolerance (e.g., associated diarrhea or upper respiratory tract infection);
7. For Nexletol, documentation that ezetimibe will be prescribed concurrently for as long as Nexletol is prescribed, unless contraindicated per Appendix F or member has a history of ezetimibe intolerance (e.g., associated diarrhea or upper respiratory tract infection);

8. Documentation of recent (within the last 60 days) LDL-C of one of the following (a or b):
  - a.  $\geq 70$  mg/dL for ASCVD;
  - b.  $\geq 100$  mg/dL for HeFH;
9. Treatment plan does not include coadministration with Repatha<sup>®</sup> or Praluent<sup>®</sup>;
10. Dose does not exceed either of the following (a or b):
  - a. Nexletol: 180 mg per day;
  - b. Nexlizet: bempedoic acid 180 mg/ezetimibe 10 mg per day.

**Approval duration: 12 months**

**B. Other diagnoses/indications**

1. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid.

**II. Continued Therapy**

**A. All Indications in Section I (must meet all):**

1. Currently receiving medication via Centene benefit or member has previously met all initial approval criteria;
2. If statin tolerant, documentation of adherence to a statin at the maximally tolerated dose;
3. For Nexletol, documentation that ezetimibe is being prescribed concurrently with Nexletol, unless a contraindication or adverse event to ezetimibe has developed;
4. Member is responding positively to therapy as evidenced by lab results within the last 3 months showing an LDL-C reduction since initiation of Nexletol or Nexlizet therapy;
5. If request is for a dose increase, new dose does not exceed either of the following (a or b):
  - a. Nexletol: 180 mg per day;
  - b. Nexlizet: bempedoic acid 180 mg/ezetimibe 10 mg per day.

**Approval duration: 12 months**

**B. Other diagnoses/indications (must meet 1 or 2):**

1. Currently receiving medication via Centene benefit and documentation supports positive response to therapy.  
**Approval duration: Duration of request or 12 months (whichever is less);** or
2. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid.

**III. Diagnoses/Indications for which coverage is NOT authorized:**

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid or evidence of coverage documents.

**IV. Appendices/General Information**

*Appendix A: Abbreviation/Acronym Key*

- |                                                  |                                            |
|--------------------------------------------------|--------------------------------------------|
| ASCVD: atherosclerotic cardiovascular disease    | LDL-C: low density lipoprotein cholesterol |
| CHD: coronary heart disease                      | SAMS: statin-associated muscle symptoms    |
| FDA: Food and Drug Administration                | TIA: transient ischemic attack             |
| HeFH: heterozygous familial hypercholesterolemia | WHO: World Health Organization             |

*Appendix B: Therapeutic Alternatives*

*This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent for all relevant lines of business and may require prior authorization.*

Drug Name	Dosing Regimen	Dose Limit/Maximum Dose
ezetimibe (Zetia <sup>®</sup> )	10 mg PO QD	10 mg/day
atorvastatin (Lipitor <sup>®</sup> )	40 mg PO QD	80 mg/day
rosuvastatin (Crestor <sup>®</sup> )	5 - 40 mg PO QD	40 mg/day

*Therapeutic alternatives are listed as Brand name<sup>®</sup> (generic) when the drug is available by brand name only and generic (Brand name<sup>®</sup>) when the drug is available by both brand and generic.*

*Appendix C: Contraindications/Boxed Warnings*

- Contraindication(s):
  - Nexletol: none reported
  - Nexlizet: known hypersensitivity to ezetimibe tablets
- Boxed warning(s): none reported

*Appendix D: Criteria for Diagnosis of HeFH*

- Dutch Lipid Clinic Network criteria for Familial Hypercholesterolemia (FH)

FH Criteria	Points	Member's Score†
<b>Family History</b>		
First-degree relative with known premature* coronary and vascular disease	1	Place highest score here (0, 1 or 2)
First-degree relative with known LDL-C level above the 95 <sup>th</sup> percentile	1	
First-degree relative with tendinous xanthomata and/or arcus cornealis	2	
Children aged < 18 years with LDL-C level above the 95 <sup>th</sup> percentile	2	

FH Criteria	Points	Member's Score†
<b>Clinical History</b>		
Patient with premature* coronary artery disease	2	Place highest score here (0, 1 or 2)
Patient with premature* cerebral or peripheral vascular disease	1	
<b>Physical Examination</b>		
Tendinous xanthomata	6	Place highest score here (0, 4 or 6)
Arcus cornealis prior to age 45 years	4	
<b>Cholesterol Levels - mg/dL (mmol/liter)</b>		
LDL-C $\geq$ 330 mg/dL ( $\geq$ 8.5)	8	Place highest score here (0, 1, 3, 5 or 8)
LDL-C 250 – 329 mg/dL (6.5 – 8.4)	5	
LDL-C 190 – 249 mg/dL (5.0 – 6.4)	3	
LDL-C 155 – 189 mg/dL (4.0 – 4.9)	1	
<b>DNA Analysis</b>		
Functional mutation in the <i>LDLR</i> , <i>apo B</i> or <i>PCSK9</i> gene	8	Place score here (0 or 8)
<b>TOTAL SCORE</b>	Definite FH: > 8	Place total score here ___

\*Premature – men < 55 years or women < 60 years

†Choose the highest score from each of the five categories and then add together for a total score. The five categories are 1) Family History, 2) Clinical History, 3) Physical Examination, 4) Cholesterol Levels, and 5) DNA Analysis.

- Simon Broome Register Group Definition of Definite FH (meets 1 and 2):
  1. One of the following (a or b):
    - a. Total cholesterol level above 7.5 mmol/l (290 mg/dl) in adults or a total cholesterol level above 6.7 mmol/l (260 mg/dl) for children under 16
    - b. LDL levels above 4.9 mmol/l (190 mg/dl) in adults (4.0 mmol/l in children) (either pre-treatment or highest on treatment)
  2. One of the following (a or b):
    - a. Tendinous xanthomas in patient or relative (parent, child, sibling, grandparent, aunt, uncle)
    - b. DNA-based evidence of an LDL receptor mutation or familial defective apo B-100
- High and Moderate Risk of ASCVD:
  - Patients with high risk of ASCVD include the following:
    - History of clinical atherosclerotic cardiovascular disease (as defined in section II)
    - Diabetes with an estimated 10-year ASCVD risk  $\geq$  7.5% for adults 40-75 years of age
    - Untreated LDL  $\geq$  190 mg/dL
  - Patients with moderate risk of ASCVD include the following:
    - Diabetes with an estimated 10-year ASCVD risk < 7.5% for adults 40-75 years of age
    - Estimated 10-year ASCVD risk  $\geq$  5% for adults 40-75 years of age

- The calculator for the 10-year ASCVD risk estimator can be found here: <http://tools.acc.org/ASCVD-Risk-Estimator-Plus/#!/calculate/estimate>. Information needed to complete the ASCVD Risk Estimator include: gender, race (white, African American, other), systolic blood pressure, history of diabetes, age, total cholesterol, HDL-cholesterol, treatment for hypertension, smoking history or status, and concurrent statin or aspirin therapy.

*Appendix E: High, Moderate, and Low Intensity Daily Statin Therapy for Adults*

<p><b>High Intensity Statin Therapy</b>  <i>Daily dose shown to lower LDL-C, on average, by approximately <math>\geq 50\%</math></i></p> <ul style="list-style-type: none"> <li>• Atorvastatin 40-80 mg</li> <li>• Rosuvastatin 20-40 mg</li> </ul>
<p><b>Moderate Intensity Statin Therapy</b>  <i>Daily dose shown to lower LDL-C, on average, by approximately 30% to 50%</i></p> <ul style="list-style-type: none"> <li>• Atorvastatin 10-20 mg</li> <li>• Fluvastatin XL 80 mg</li> <li>• Fluvastatin 40 mg BID</li> <li>• Lovastatin 40 mg</li> <li>• Pitavastatin 1-4 mg</li> <li>• Pravastatin 40-80 mg</li> <li>• Rosuvastatin 5-10 mg</li> <li>• Simvastatin 20-40 mg</li> </ul>
<p><b>Low Intensity Statin Therapy</b>  <i>Daily dose shown to lower LDL-C, on average, by <math>&lt; 30\%</math></i></p> <ul style="list-style-type: none"> <li>• Simvastatin 10 mg</li> <li>• Pravastatin 10-20 mg</li> <li>• Lovastatin 20 mg</li> <li>• Fluvastatin 20-40 mg</li> </ul>

*Appendix F: Statin and Ezetimibe Contraindications*

<p><b>Statins</b></p> <ul style="list-style-type: none"> <li>• Decompensated liver disease (development of jaundice, ascites, variceal bleeding, encephalopathy)</li> <li>• Laboratory-confirmed acute liver injury or rhabdomyolysis resulting from statin treatment</li> <li>• Pregnancy*, actively trying to become pregnant, or nursing</li> <li>• Immune-mediated hypersensitivity to the HMG-CoA reductase inhibitor drug class (statins) as evidenced by an allergic reaction occurring with at least TWO different statins</li> </ul>
<p><b>Ezetimibe</b></p> <ul style="list-style-type: none"> <li>• Moderate or severe hepatic impairment [Child-Pugh classes B and C]</li> <li>• Hypersensitivity to ezetimibe (e.g., anaphylaxis, angioedema, rash, urticaria)</li> </ul>

*\*In July 2021, the FDA requested removal of the contraindication against use of statins in pregnant women. Because the benefits of statins may include prevention of serious or potentially fatal events in a small group of very high-risk pregnant patients, contraindicating these drugs in all pregnant women is not appropriate.*

<https://www.fda.gov/safety/medical-product-safety-information/statins-drug-safety-communication-fda-requests-removal-strongest-warning-against-using-cholesterol>

*Appendix G: Statin Risk Factors*

<b>Statin Risk Factors</b>
<ul style="list-style-type: none"> <li>• Multiple or serious comorbidities, including impaired renal or hepatic function</li> <li>• Unexplained alanine transaminase (ALT) elevations &gt; 3 times upper limit of normal, or active liver disease</li> <li>• Concomitant use of drugs adversely affecting statin metabolism</li> <li>• Age &gt; 75 years, or history of hemorrhagic stroke</li> <li>• Asian ancestry</li> </ul>

*Appendix H: General Information*

- Patients should remain on concomitant therapy with a statin if tolerated due to the established long term cardiovascular benefits.
- The diagnosis of SAMS is often on the basis of clinical criteria. Typical SAMS include muscle pain and aching (myalgia), cramps, and weakness. Symptoms are usually bilateral and involve large muscle groups, including the thigh, buttock, back, and shoulder girdle musculature. In contrast, cramping is usually unilateral and may involve small muscles of the hands and feet. Symptoms may be more frequent in physically active patients. Symptoms often appear early after starting statin therapy or after an increase in dose and usually resolve or start to dissipate within weeks after cessation of therapy, although it may take several months for symptoms to totally resolve. Persistence of symptoms for more than 2 months after drug cessation should prompt a search for other causes or for underlying muscle disease possibly provoked by statin therapy. The reappearance of symptoms with statin rechallenge and their disappearance with drug cessation offers the best evidence that the symptoms are truly SAMS.
- Pravastatin, fluvastatin, and rosuvastatin are hydrophilic statins which have been reported to confer fewer adverse drug reactions than lipophilic statins.

**V. Dosage and Administration**

<b>Indication</b>	<b>Dosing Regimen</b>	<b>Maximum Dose</b>
ASCVD, HeFH	One tablet PO QD	One tablet/day

**VI. Product Availability**

<b>Drug Name</b>	<b>Availability</b>
Bempedoic acid (Nexletol)	Tablet: 180 mg
Bempedoic acid/ezetimibe (Nexlizet)	Tablet: bempedoic acid 180 mg/ezetimibe 10 mg

**VII. References**

1. Nexletol Prescribing Information. Ann Arbor, MI: Esperion Therapeutics, Inc.; February 2020. Available at: [www.Nexletolhcp.com](http://www.Nexletolhcp.com). Accessed October 1, 2021.
2. Nexlizet Prescribing Information. Ann Arbor, MI: Esperion Therapeutics, Inc.; February 2020. Available at: <https://pi.esperion.com/nexlizet/nexlizet-pi.pdf>. Accessed October 1, 2021.

3. Grundy SM, Stone NJ, Bailey AL, et al. 2018 ACC/AHA/AACVPR/AAPA/ABC/ACPM/ADA/AGS/APhA/ASPC/NLA/PCNA guideline on the management of blood cholesterol: A report of the American College of Cardiology Foundation/American Heart Association Task Force on Clinical Practice Guidelines. J Am Coll Cardiol 2018;Nov 10:[Epub ahead of print].
4. Backes JM, Ruisinger JF, Gibson CA, et al. Statin-associated muscle symptoms—managing the highly intolerant. J Clin Lipidol. 2017;11:24-33. Available at: <https://www.acc.org/latest-in-cardiology/ten-points-to-remember/2017/05/03/10/43/statin-associated-muscle-symptoms>. Accessed October 1, 2021.
5. Goldberg AC, Letier LA, Stroes ESG, et al. Effect of bempedoic acid vs placebo added to maximally tolerated statins on low-density lipoprotein cholesterol in patients at high risk for cardiovascular disease – the CLEAR Wisdom randomized clinical trial. JAMA 2019;322(18):1780-8.
6. Ray KK, Bays HE, Catapano AL, et al. Safety and efficacy of bempedoic acid to reduce LDL cholesterol. N Engl J Med 2019;380:1022-32.
7. Ballantyne CM, Laufs U, Ray KK, et al. Bempedoic acid plus ezetimibe fixed-dose combination in patients with hypercholesterolemia and high CVD risk treated with maximally tolerated statin therapy. Eur J Prev Cardiol 2019 Jul 29 [online ahead of print]; DOI: 10.1177/2047487319864671.
8. DRUGDEX<sup>®</sup> System [Internet database]. Greenwood Village, Colo: Thomson Healthcare. Updated periodically. Accessed October 1, 2021.
9. Institute for Clinical and Economic Review. Bempedoic acid and inclisiran for patients with heterozygous familial hypercholesterolemia and for secondary prevention of ASCVD: Effectiveness and value. Final evidence report published March 2, 2021. Available at: <https://icer.org/assessment/high-cholesterol-2021>. Accessed October 1, 2021.

Reviews, Revisions, and Approvals	Date	P&T Approval Date
Policy created	04.07.20	08.20
1Q 2021 annual review: no significant changes; references to HIM.PHAR.21 revised to HIM.PA.154; references reviewed and updated.	11.02.20	02.21
1Q 2022 annual review: no significant changes; references reviewed and updated.	10.01.21	02.22

**Important Reminder**

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. “Health Plan” means a health



plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan's affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

**Note:**

**For Medicaid members**, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

©2020 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise

## CLINICAL POLICY

### Bempedoic Acid, Bempedoic Acid/Ezetimibe



published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene® and Centene Corporation® are registered trademarks exclusively owned by Centene Corporation.